

Authors

Mario Angelico

*Liver and Kidney Transplant Unit
Tor Vergata University
Rome, Italy*

Marina Berenguer

*Gastroenterology Service
Hepatology Unit
CIBERehd
Hospital La Fe
Valencia, Spain*

Andres F. Carrion

*Department of Medicine
University of Miami
Miller School of Medicine
Miami, FL, USA*

José A. Carrión

*Liver Unit
Institut de Malalties Digestives
Hospital Clinic, IDIBAPS, CIBERehd
Barcelona, Spain*

Chan Y. Chung

*Division of Gastroenterology
University of California
San Francisco
San Francisco, CA, USA*

Gonzalo Crespo

*Liver Unit
Institut de Malalties Digestives
Hospital Clinic, IDIBAPS, CIBERehd
Barcelona, Spain*

Fabrizio Fabrizi

*Division of Nephrology and Dialysis
Maggiore Hospital
IRCCS Foundation
Milan, Italy*

Hélène Fontaine

*Paris Descartes University
Hepatology Unit
AP-HP Hôpital Cochin
Paris, France*

Xavier Forns

*Liver Unit
Institut de Malalties Digestives
Hospital Clinic, IDIBAPS, CIBERehd
Barcelona, Spain*

Juan F. Gallegos-Orozco

*Division of Gastroenterology
Mayo Clinic
Scottsdale, AZ, USA*

Tommaso Maria Manzia

*Liver and Kidney Transplant Unit
Tor Vergata University
Rome, Italy*

Paul Martin

*Division of Hepatology
University of Miami
Miller School of Medicine
Miami, FL, USA*

Stanislas Pol

*Paris Descartes University
Hepatology Unit
AP-HP Hôpital Cochin
Paris, France*

Bruno Roche

*Hepato Biliary Center
AP-HP Hôpital Paul Brousse
University Paris-Sud
Villejuif, France*

Didier Samuel

*Hepato Biliary Center
AP-HP Hôpital Paul Brousse
University Paris-Sud
Villejuif, France*

Norah A. Terrault

*Division of Gastroenterology
University of California
San Francisco
San Francisco, CA, USA*

Giuseppe Tisone

*Liver and Kidney Transplant Unit
Tor Vergata University
Rome, Italy*

Anais Vallet-Pichard

*Paris Descartes University
Hepatology Unit
AP-HP Hôpital Cochin
Paris, France*

Hugo E. Vargas

*Division of Hepatology
Mayo Clinic
Phoenix, AZ, USA*